



A regularized vortex-particle mesh method for large eddy simulation

Spietz, Henrik Juul; Walther, Jens Honore; Hejlesen, Mads Mølholm

Publication date:
2017

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Spietz, H. J., Walther, J. H., & Hejlesen, M. M. (2017). *A regularized vortex-particle mesh method for large eddy simulation*. Abstract from 70th Annual Meeting of the American Physical Society Division of Fluid Dynamics (DFD17), Denver, Colorado, United States.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Abstract Submitted
for the DFD17 Meeting of
The American Physical Society

Sorting Category: 37.8 (C)

A regularized vortex-particle mesh method for large eddy simulation H. J. SPIETZ, Technical University of Denmark, J. H. WALTHER, Technical University of Denmark and ETH Zurich, M. M. HEJLESEN, Technical University of Denmark — We present recent developments of the remeshed vortex particle-mesh method for simulating incompressible fluid flow. The presented method relies on a parallel higher-order FFT based solver for the Poisson equation. Arbitrary high order is achieved through regularization of singular Green's function solutions to the Poisson equation and recently we have derived novel high order solutions for a mixture of open and periodic domains. With this approach the simulated variables may formally be viewed as the approximate solution to the filtered Navier Stokes equations, hence we use the method for Large Eddy Simulation by including a dynamic subfilter-scale model based on test-filters compatible with the aforementioned regularization functions. Further the subfilter-scale model uses Lagrangian averaging, which is a natural candidate in light of the Lagrangian nature of vortex particle methods. A multiresolution variation of the method is applied to simulate the benchmark problem of the flow past a square cylinder at $Re = 22000$ and the obtained results are compared to results from the literature.

☒ Prefer Oral Session
☐ Prefer Poster Session

Henrik J. Spietz
hejusp@mek.dtu.dk
Technical University of Denmark

Date submitted: 28 Jul 2017

Electronic form version 1.4